

469253-continued

T 3707. 'PMT - 389'. United States. Collected 1958. Van Horn, Texas. Adapted to medium and fine textured soils of western Edwards Plateau, Rio Grande Plain, and Trans Pecos area of west Texas. Seed adheres to grama seed and feeds readily through seeding equipment. Perennial Herb. Cultivar. Seed.

469254. *Eragrostis superba* Peyr. (Poaceae) Wilman lovegrass.

From United States. Knox City Plant Materials Center, Soil Conservation Service, USDA, Los Lunas, Mexico.
Received April 1982.

T 8237. South Africa. Composite of PI 295704, PI 299959, BN 14836, and BN 16087. Adapted to sandy and heavy, black soil. Limited to areas where minimum temperature is 10 deg. F. Used for forage, wildlife and erosion control on desert grassland. More palatable than weeping or Lehman lovegrass. Preferred to Rhodesgrass and blue panicum in irrigated pastures. Perennial Herb. Cultivated. Seed.

469255 TO 469257. *Leucaena leucocephala* (Lam.) de Wit (Fabaceae) Leadtree.

From Brazil. C. C. Shock, Department of Agronomy & Range Science, University of California, Davis, CA.
Received May 1982.

469255. Peruvian. Brazil. Sao Paulo. Instituto de Pesquisas. IRI 2984. Originally from Australia. Cultivated. Seed.

469256. Brazil. Sao Paulo. Instituto de Pesquisas. IRI 3155. Collected by C. C. Shock. Single plant growing in Makueni guineagrass. Cultivated. Seed.

469257. Brazil. Sao Paulo. Instituto de Pesquisas. IRI 3164. Brasilandia, Mato Grosso do Sul. Single plant selection from crop of Peruvian. Cultivated. Seed.

469258 TO 469260. *Panicum maximum* Jacq. (Poaceae) Guineagrass.

From Brazil. C. C. Shock, Department of Agronomy & Range Science, University of California, Davis, CA.
Received May 1982.

469258. Brazil. Selection A from CPI 37910. Cultivated. Seed.

469259. Brazil. Selection B from CPI 37910. Cultivated. Seed.

469260. Brazil. Selection C from CPI 37910. Cultivated. Seed.